



King's Research Portal

DOI:

[10.1111/dmcn.13707](https://doi.org/10.1111/dmcn.13707)

Document Version

Peer reviewed version

[Link to publication record in King's Research Portal](#)

Citation for published version (APA):

Shortland, A. (2018). Muscle tone is not a well-defined term. *Developmental Medicine and Child Neurology*, 60(7). <https://doi.org/10.1111/dmcn.13707>

Citing this paper

Please note that where the full-text provided on King's Research Portal is the Author Accepted Manuscript or Post-Print version this may differ from the final Published version. If citing, it is advised that you check and use the publisher's definitive version for pagination, volume/issue, and date of publication details. And where the final published version is provided on the Research Portal, if citing you are again advised to check the publisher's website for any subsequent corrections.

General rights

Copyright and moral rights for the publications made accessible in the Research Portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognize and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the Research Portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the Research Portal

Take down policy

If you believe that this document breaches copyright please contact librarypure@kcl.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.

[1 Commentary; Adam.Shortland@gstt.nhs.uk]

Muscle tone is not a well-defined term

ADAM PETER SHORTLAND

Guy's & St Thomas' Foundation Hospital Trust – One Small Step Gait
Laboratory, London, UK.

School of Biomedical Engineering and Imaging Sciences, King's College
London, UK.

doi: 10.1111/dmcn.

This commentary is on the systematic review by Goo et al. To view this
paper visit <https://doi.org/10.1111/dmcn>.

'When I use a word ... it means just what I choose it to mean — neither
more nor less.'

Lewis Carroll's Humpty Dumpty might well have been thinking of the word
'tone' while he was philosophising with Alice. Tone is a commonly-used
term most notable for its ambiguity. Definitions of tone (e.g. Sanger et
al.¹) contain phrases and words which are themselves ill-defined,
rendering them difficult to interpret or measure. For example, Sanger et
al. define passive tone as 'resistance to passive stretch while a patient is
attempting to maintain a relaxed state of muscle activity,' but what is

meant *precisely* by resistance to stretch? Any definition that contains words or phrases that are ambiguous will always give the individual clinician a little room for manoeuvre to create their own internalized, even subconscious, version of the concept.

You would think that having a shared concept which was a little different in each clinician's mind would be in some ways progress-limiting, frustrating or worse, counter-productive. However, having a loosely-defined concept, and accepting that other clinicians' internalized versions are not quite the same as your own, allows us to communicate with each other in broad terms without getting hung up on the detail. So, if a colleague says to me, 'I saw a child with excessive tone this morning,' I have a mental image of what that means, what that child might 'look' like. Useful, if we both understand the limits of the shared ambiguous concept.

Problems occur when we take concepts like tone and we try to measure something about them, like their psychometric properties. In their systematic review, Goo et al. have bravely attempted to do just that, but have they achieved their objective?² In fact, they have analysed the properties of the components of developmental scales that could only be very loosely associated with our ambiguous concept of tone. They claim, for example, that 21 components from the numerous developmental scales they reviewed were consistent with Sanger et al.'s definition of passive tone. Actually, they selected items that were based on observation, palpation, passive range of motion, *or* resistance to

stretch. These terms are not all consistent with Sanger et al.'s definition, insofar as passive range of motion is not the same thing as resistance to stretch. One might also argue that each of these terms have rather ambiguous meanings themselves and therefore might have a very different representation in each assessor's mind. No wonder then that most of the assessments of tone featured in this review had poor to fair validity, and that only one test achieved moderate levels agreement between raters. Perhaps even more concerning was the poor intrarater reliability (assessors didn't agree with themselves on retesting) which suggest that either the status of the participants is very variable, the measurements intrinsically variable, or that the concept in the assessor's head changes. In any case, the components of the developmental scales reviewed by Goo et al.² are likely to have limited value in charting neurological development and thus predicting developmental problems.

REFERENCES

1. Sanger TD, Delgado MR, Gaebler-Spira D, Hallett M, Mink JW ; Task Force on Childhood Motor Disorders. Classification and definition of disorders causing hypertonia in childhood. *Pediatrics* 2003; **111**: e89–97.
2. Goo M, Tucker K, Johnson LM. Muscle tone assessments for children aged 0 to 12 years: a systematic review. *Dev Med Child Neurol* 2018; <https://doi.org/10.1111/dmcn>.